

Serial No.: 10/691,157

Confirmation No.: 6536

Filed: 22 October 2003

For: USE OF COLOSTRININ, CONSTITUENT PEPTIDES THEREOF, AND ANALOGS THEREOF AS  
MODULATORS OF INTRACELLULAR SIGNALING MOLECULES

---

**Amendments to the Specification**

Please replace the paragraph beginning at page 12, line 15, with the following amended paragraph.

As stated above, active analogs of colostrinin and its constituent peptides include polypeptides having structural similarity. Structural similarity is generally determined by aligning the residues of the two amino acid sequences to optimize the number of identical amino acids along the lengths of their sequences; gaps in either or both sequences are permitted in making the alignment in order to optimize the number of identical amino acids, although the amino acids in each sequence must nonetheless remain in their proper order. Preferably, two amino acid sequences are compared using the Blastp program, version 2.0.9, of the BLAST 2 search algorithm, available on the world wide web at <http://www.ncbi.nlm.nih.gov/gorf/bl2.html>. Preferably, the default values for all BLAST 2 search parameters are used, including matrix = BLOSUM62; open gap penalty = 11, extension gap penalty = 1, gap x\_dropoff = 50, expect = 10, wordsize = 3, and filter on. In the comparison of two amino acid sequences using the BLAST search algorithm, structural similarity is referred to as "identity." Preferably, an active analog of colostrinin or its constituent peptides has a structural similarity to colostrinin or one or more of its constituent peptides (preferably, one of SEQ ID NOs:1-34) of at least about 70% identity, more preferably, at least about 80% identity, and most preferably, at least about 90% identity.